

# UNCLASSIFIED

AD NUMBER
AD270145
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 04 SEP 1961. Other requests shall be referred to Army Medical Research and Development Command, Aberdeen Proving Ground, MD.
AUTHORITY
USAMA ltr, 26 Feb 1970

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

---

AD 270 145

*Reproduced  
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA



---

UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

11/

# US ARMY MEDICAL RESEARCH LABORATORY

FORT KNOX, KENTUCKY

REPORT NO. 504

CESIUM-137 RETENTION AND DISTRIBUTION IN  
X-IRRADIATED RATS



J. G. Kereiakes, Ph. D.  
Capt D. D. Ulmer, MC  
A. T. Krebs, Ph.D.  
T. D. Sterling

270 145

AD NO.

ASTIA FILE COPY

270 145



UNITED STATES ARMY

MEDICAL RESEARCH AND DEVELOPMENT COMMAND 4 September 1961

NO OTS

Report Submitted 1 September 1961

Authors

J. G. Kereiakes, Ph.D. *	Chief, Acute Irradiation Branch Radiobiology Division
Captain D. D. Ulmer, MC** (M. D. )	Acute Irradiation Branch Radiobiology Division
A. T. Krebs, Ph.D.	Director Radiobiology Division
T. D. Sterling	Biometrics Division College of Medicine University of Cincinnati Cincinnati, Ohio

\*Now at the University of Cincinnati, Cincinnati, Ohio.

\*\*Now at the Peter Bent Brigham Hospital, Boston, Massachusetts.

Qualified requestors may obtain copies of this report from ASTIA.

Foreign announcement and dissemination of this report by ASTIA is limited.

AD	Accession No.	UNCLASSIFIED	AD	Accession No.	UNCLASSIFIED
US Army Medical Research Lab., Ft. Knox, Ky. CESIUM-137 RETENTION AND DISTRIBUTION IN X- IRRADIATED RATS. J. G. Kretzschmar, D. D. Ulmer, A. T. Krebs, and T. D. Sterling Report No. 504, 4 Sep 61, 12 pp & ill. 4 illus. 3 tables. Project No. 6X64-14-001. Unclassified Report		1. Radiobiology 2. Isotope 3. Cesium-137	US Army Medical Research Lab., Ft. Knox, Ky. CESIUM-137 RETENTION AND DISTRIBUTION IN X- IRRADIATED RATS. J. G. Kretzschmar, D. D. Ulmer, A. T. Krebs, and T. D. Sterling Report No. 504, 4 Sep 61, 12 pp & ill. 4 illus. 3 tables. Project No. 6X64-14-001. Unclassified Report		1. Radiobiology 2. Isotope 3. Cesium-137
<p>The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1 <math>\mu</math>c cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5 <math>\mu</math>c cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.</p>			<p>The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1 <math>\mu</math>c cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5 <math>\mu</math>c cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.</p>		
AD	Accession No.	UNCLASSIFIED	AD	Accession No.	UNCLASSIFIED
US Army Medical Research Lab., Ft. Knox, Ky. CESIUM-137 RETENTION AND DISTRIBUTION IN X- IRRADIATED RATS. J. G. Kretzschmar, D. D. Ulmer, A. T. Krebs, and T. D. Sterling Report No. 504, 4 Sep 61, 12 pp & ill. 4 illus. 3 tables. Project No. 6X64-14-001. Unclassified Report		1. Radiobiology 2. Isotope 3. Cesium-137	US Army Medical Research Lab., Ft. Knox, Ky. CESIUM-137 RETENTION AND DISTRIBUTION IN X- IRRADIATED RATS. J. G. Kretzschmar, D. D. Ulmer, A. T. Krebs, and T. D. Sterling Report No. 504, 4 Sep 61, 12 pp & ill. 4 illus. 3 tables. Project No. 6X64-14-001. Unclassified Report		1. Radiobiology 2. Isotope 3. Cesium-137
<p>The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1 <math>\mu</math>c cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5 <math>\mu</math>c cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.</p>			<p>The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1 <math>\mu</math>c cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5 <math>\mu</math>c cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.</p>		

REPORT NO. 504

CESIUM-137 RETENTION AND DISTRIBUTION IN  
X-IRRADIATED RATS

J. G. Kereiakes, Ph.D.  
Capt D. D. Ulmer, MC  
A. T. Krebs, Ph.D.  
T. D. Sterling

Radiobiology Division  
US ARMY MEDICAL RESEARCH LABORATORY  
Fort Knox, Kentucky

4 September 1961

Effects of Low, Medium, and Massive Doses  
Task 01  
Biological and Medical Aspects of Ionizing Radiation  
USAMRL Project No. 6X64-14-001

ABSTRACT

**CESIUM-137 RETENTION AND DISTRIBUTION IN  
X-IRRADIATED RATS**

OBJECT

To study the influence of external ionizing radiation on the retention and distribution of the long-lived fission-product, cesium-137, in the rat.

RESULTS

Immediately following a total body exposure of 300, 600, and 900 r of 250 kv X-rays, rats were injected intraperitoneally with 1  $\mu$ c solution of cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium over the three-day period following irradiation exposure than did the control group. In a separate tissue distribution study, the largest part of the injected cesium dose was found in the muscle. Cesium was also concentrated to a smaller extent in the small intestine, liver, bone, and large intestine. The irradiated rats retained less cesium-137, expressed as mean per cent of injected dose, in muscle, bone, blood, small intestine, and spleen; whereas more cesium was held in the stomach and liver. The cesium content of whole organs was reflected, with the exception of stomach and spleen, to the same extent when considered on an organ unit-weight basis.

CONCLUSIONS

An analysis of the data indicated that the increased cesium excretion in the X-irradiated rats was primarily related to the greater urine volume (diuresis accompanied by increased water intake) in these rats resulting from radiation exposure. Increased urine excretion of cesium probably accounted for most of the lower values observed for cesium in the organs of the irradiated rats. The gastrointestinal concentrations of cesium-137 are explainable on the basis of gastric retention and delayed intestinal motility characteristic of the rat species following irradiation exposure. The liver concentration may be related to its corresponding weight increase following irradiation exposure; whereas the

spleen appeared to lose weight at a greater rate than it lost cesium. The failure of bone in the irradiated rats to take up cesium to the same extent as the control rats may be due to decreased blood concentration of cesium in these rats or it may reflect direct irradiation effects on the bone.

#### RECOMMENDATIONS

The cesium-137 concentration in the bone of X-irradiated rats should be studied for various radiation doses and for various times following irradiation exposure. The cesium-137 distribution in rats given a diuretic drug should be studied and compared with the findings reported in this study for X-irradiation exposure.

APPROVED:

A. T. Krebs  
A. T. KREBS, Ph. D.  
Director, Radiobiology Division

APPROVED:

Floyd A. Odell  
FLOYD A. ODELL, Ph. D.  
Technical Director of Research

APPROVED:

Sven A. Bach  
SVEN A. BACH  
Colonel, MC  
Commanding

# CESIUM-137 RETENTION AND DISTRIBUTION IN X-IRRADIATED RATS

## I. INTRODUCTION

Although the animal and human metabolism of fission-products has been extensively explored, the influence of external ionizing radiation on the uptake, distribution, and excretion of such elements has received comparatively little attention. Since both environmental exposure and clinical toxicity of these two sources of radiation may be interdependent, an experimental investigation of their combined short term effects in mammals has been undertaken.

Ulmer, Perkins, and Kerciakes (1) studied the distribution of iodine-131 in rats 24 hours after whole-body X-irradiation and found marked alterations compared to the isotope distribution in non-irradiated control animals. A reduction of 20 per cent in urinary and fecal loss of radioiodine occurred post-irradiation. As a consequence, increased radioactivity appeared in the blood, most of the organs, and particularly in the contents of the stomach and the pelt.

The present report is concerned with more detailed investigations of the retention and distribution of the longer-lived fission product, cesium-137, in rat tissue following whole-body radiation exposure.

## II. EXPERIMENTAL

For exposure to whole-body X-irradiation the rats were placed two at a time in a shallow lucite cage. X-rays were delivered by a General Electric Maxitron Unit operated at 250 kvp, 30 ma, 4.75 mm Be inherent filtration, 1 mm Al + 0.5 mm Cu added filtration, HVL 1.1 mm Cu. The dose measured in air with a 250 r Victoreen Ionization Chamber was 140 r/min at a target-midline rat distance of 50 cm.

### Cesium Retention

Male Sprague-Dawley rats weighing 160-250 grams were divided into the following groups: Cesium-137 injected only (controls), and cesium-137 with 300 r, 600 r, or 900 r whole-body radiation exposure.

Immediately following X-ray exposure the rats were injected intraperitoneally with 1 ml of a neutral aqueous solution containing 1  $\mu$ c of cesium-137 (chemical form - CsCl). An aliquot of the solution was

set aside to be used as a standard. The rats were housed individually in metabolism cages to allow both separation and quantitative collection of urine and feces and accurate measurement of the water ingested. The rats were starved but were permitted water ad libitum. Control rats (those receiving cesium-137 but not irradiated) were handled in the same manner as the irradiated rats. Ten minutes after injection of the isotope the animal was placed in a glass tube (1-3/4 inches in diameter by 8 inches long) having holes at one end for air and a rubber stopper at the other end. The rat and tube were placed inside the well of a Nucleonic Corporation of America Supersensitive Large-Well Geiger-Counter (Fig. 1). The well in this counter measures 2 inches in diameter by 12 inches in length. This well-counter consists of multicylindrical elements with mesh walls concentrically arranged around the well. Its pulse size and shape are the same as the ordinary Geiger-Müller pulse. However, the above element arrangement overcomes to some extent the recognized limitations of the ordinary Geiger-counter, e. g., namely the insensitivity to gamma rays, long resolution time, and short life. The well-counter was connected to a Radiation Counter Laboratory Scaler and was shielded all around with 1-1/2 inches of lead.

The whole-body count measured 10 minutes after injection of cesium served as the control for each animal (i. e., the "100 per cent whole-body count"). Whole-body counts on each rat were repeated at 7 hours, and at 1, 2, and 3 days following injection. Total water ingestion, urine volume, and feces weight were measured each 24 hours. Aliquots of urine and digested feces (after digestion in nitric acid) were placed in test tubes and counted with a Radiation Counter Laboratory Well-Type Scintillation counter having a NaI (Tl) crystal.

#### Cesium Distribution

Male Sprague-Dawley rats weighing 170-220 grams were selected from litter mates taken after weaning and housed four to a cage until they attained the desired weight. Immediately after exposure to 900 r irradiation under the conditions described above, the two irradiated rats as well as two control (sham-irradiated) litter mates were injected intraperitoneally with a 1 ml saline solution containing 5  $\mu$ c cesium-137. At the time of injection an aliquot of the cesium solution was set aside for use as a standard. The animals were then separately housed in metabolism cages. Food was withheld but water was given ad libitum.

Twenty-four hours after X-ray exposure the rats were lightly anesthetized with ether and killed by exsanguination from the abdominal

aorta. The major internal organs were removed intact. A muscle sample was taken from the left thigh while the right gonad and right femur were utilized for analysis. All organs were stripped of extraneous connective tissue, washed carefully, blotted dry, and weighed. The contents of the stomach, small intestine, and large intestine were included in the analyses of these organs; feces were added to the contents of the large intestine. Whole organs or 1 to 2 gram portions were weighed and taken for analysis. All samples were wet ashed in hot concentrated nitric acid and diluted as necessary for counting. Samples were counted in a volume of 3 ml in the well-type scintillation counter described above.

For final calculations, muscle was considered to constitute 45 per cent of whole-body weight and blood volume 5.3 ml per 100 gram body weight; bone weight was estimated from standard tables (2).

### III. RESULTS

#### Cesium Retention

The results of the retention study are shown in Tables 1 and 2. Rats receiving cesium-137 only will be referred to as the control group. The body weight of the control group (food withheld) decreased to 77.7 per cent of the initial body weight in 3 days; whereas the body weight of the irradiated rats decreased to 74.3 to 77.2 per cent of their initial weight during this time interval.

Exposure to ionizing radiation significantly alters the retention of cesium-137 by the body of the rat. The whole-body content of this isotope at 7, 24, 48, and 72 hours after exposure to irradiation of 300 r to 900 r is shown graphically in Figure 2. Uniformly, and from the time of earliest measurement, the irradiated animal retained less cesium-137 than did the corresponding non-irradiated controls. Moreover, this response to irradiation was of progressively greater significance with time, being most marked on the final day of observation. The effect was not directly proportional to radiation dose, however, being maximal in those animals exposed to 600 r.

As a necessary corollary, exposure to ionizing radiation should significantly increase the rate of elimination of cesium in rats. This is demonstrated by measurement of urinary cesium-137 radioactivity (Fig. 3). In all instances, urinary excretion of the isotope is increased subsequent to X-irradiation. Excretion by this route was maximal on

the first day after exposure to X-ray (up to 12 per cent of the injected isotope) and fell off rapidly thereafter. The quantity of radioactivity in the feces was small, amounting to less than 1 per cent of the injected dose of cesium for both groups and appeared to be proportional to the volume of fecal material (Tables 1 and 2).

#### Cesium Distribution

The data are expressed as 1) mean per cent of injected dose per total organ, and 2) mean concentration of radioactivity (counts per second per gram of sample divided by counts per second injected per gram of body weight) (3). The latter values are calculated to compensate for changes in organ weight and variations in body weight. These results are given in Table 3 along with calculations for the "t" test and probability values.

In both control and X-irradiated animals the greater portion of the injected isotope was found in muscle. Analyses of the other tissues indicated that cesium was also concentrated, but, to a lesser extent, in the small intestine, liver, bone, and large intestine. The data for all the tissues studied are shown graphically in Figure 4.

Irradiated rats retained less cesium-137 in the muscle than did control rats, probably reflecting increased cesium excretion into the urine. As in the retention study the total urine volume was uniformly greater in the irradiated rats and was accompanied by increased content of cesium-137. There was a correspondingly greater ingestion of water in irradiated animals. The blood concentration of cesium was significantly higher in the control rats. There was no significant difference between the two groups in the 24 hour cesium content of gonads, heart, kidneys, and large intestine. Bone, small intestine, and spleen retained less cesium in the irradiated rats, while the liver and stomach contained greater concentrations of the isotope.

The distribution of cesium was, in general, unchanged when calculated on a unit weight basis. Two exceptions were the spleen and the stomach plus contents. On a unit weight basis, the spleen retained more and the stomach less cesium in the irradiated rats.

#### IV. DISCUSSION

Cesium-137 has a half-life of 30 years and emits 1.17 Mev (8 per cent) and 0.51 Mev (92 per cent) beta particles followed by 0.662 Mev

gamma rays. As one of the alkali metals, it would be predicted that the organ distribution of cesium should resemble that of potassium. In this regard excretion and distribution studies with the fission products, cesium-135 and cesium-138, were reported by Hamilton (4), who found a relatively high accumulation in the soft tissues, especially muscle, and a low uptake in the skeleton. Absorption from the intestinal tract was found to be 100 per cent after ingestion with about 50 per cent being excreted within 10 days.

Hood and Comar (5, 6) studied the metabolism of cesium-137 in the rat. Cesium was widely distributed in the body. Skeletal accumulation, represented by isotope concentration in the femur, was found to be quite small. The greatest accumulation and most tenacious retention was in muscle tissue, although this tissue accumulated cesium at a slightly lower rate over the first week than did liver, spleen, and kidney. Blood always had the lowest concentration, so the tissues may have taken up cesium against a concentration gradient.

Ballou and Thompson (7) investigated the effect of acute and chronic administration of cesium-137 in rats. The experiments on acute administration of the element were of particular interest to the present discussion. After a single dose the retention of cesium-137 was followed in eleven organs and tissues for a period of 200 days. Muscle was again found to be the critical organ.

The present study both confirms results in the distribution of cesium reported by other workers and extends these data to encompass the effect of the added stress of external whole-body irradiation exposure. The results clearly indicate a greater excretion of cesium-137 in the irradiated rats, quite probably related to the increased urinary volume in these animals. Hence a major increase in the excretion of this isotope apparently may be brought about by inducing diuresis. To ascertain if, indeed, such urinary losses of cesium in irradiated animals exceeded the control values primarily as a function of increased urinary volume, the observations for the 3 days were pooled and normalized by assuming that the number of counts per urine sample was a constant fraction of the urine volume. The difference between the observed urine counts and those predicted on the basis of the changes in urinary volume were then obtained. There was no significant difference between the observed and predicted urine counts, strongly suggesting that the urinary losses of cesium in the irradiated animals and increased urinary volume are causally related phenomena.

The water metabolism in the rat following X-irradiation has been studied by Edelmann (8, 9). Both ingestion and excretion of water are increased on the first day after irradiation and are proportional to log dose, at least in the range of 75 r to 600 r. If the fluid intake was restricted for 24 hours immediately following irradiation, urinary volume still increased. Hence Edelmann felt that diuresis preceded and may, in fact, be the cause of polydipsia. Serum taken from the irradiated rats within 24 hours after exposure to irradiation and assayed for an "antidiuretic substance," probably the antidiuretic factor of the posterior lobe of the pituitary gland, was found to contain smaller concentrations of this substance than did the serum of unirradiated controls. At least the first period of diuresis, therefore, coincides with a decreased amount of circulating antidiuretic substance. It would appear that cesium in body water would be markedly affected by this diuresis and the concomitantly increased water intake contributing, in a great part, to the differences in the excretion of the element.

If the tissue deposition of cesium is examined further, several interesting features are noted. In general, the organs of irradiated animals contained less cesium, as would be predicted from the greater urinary losses in this group (Table 3). The increased gastrointestinal concentration of cesium, probably due to gastric retention and delayed intestinal motility (10, 11), may also influence the distribution pattern. Although there is more total cesium in the stomach of irradiated animals than in the control group, it is more dilute and accounts for the lesser cesium concentration in the adjoining small intestine. The variations in concentrations of cesium in the liver and spleen are most likely related to weight changes in these organs subsequent to irradiation. The liver weight increased by 24 per cent at 24 hours in the irradiated rats. It has been postulated that such a post-irradiation increase in liver size is due to an increase in glycogen storage (12). The mean weight of the spleens in the irradiated rats decreased by about 50 per cent at 24 hours. It would appear that the spleen in these animals lost weight at a greater rate than it lost cesium.

Of considerable interest is the striking failure of bone in the irradiated animal to take up cesium at 24 hours to the extent which occurs in the control animal. The concentrations in bone may be due to decreased blood concentration of cesium in these rats or it may reflect direct irradiation effects on bone. A further detailed study of this phenomenon is warranted.

#### V. CONCLUSIONS

An analysis of the data indicated that the increased cesium excretion in the X-irradiated rats was primarily related to the greater urine

volume (diuresis accompanied by increased water intake) in these rats resulting from radiation exposure. Increased urine excretion of cesium probably accounted for most of the lower values observed for cesium in the organs of the irradiated rats. The gastrointestinal concentrations of cesium-137 are explainable on the basis of gastric retention and delayed intestinal motility characteristic of the rat species following irradiation exposure. The liver concentration may be related to its corresponding weight increase following irradiation exposure; whereas the spleen appeared to lose weight at a greater rate than it lost cesium. The failure of bone in the irradiated rats to take up cesium to the same extent as the control rats may be due to decreased blood concentration of cesium in these rats or it may reflect direct irradiation effects on the bone.

#### VI. RECOMMENDATIONS

The cesium-137 concentration in the bone of X-irradiated rats should be studied for various radiation doses and for various times following irradiation exposure.

The cesium-137 distribution in rats given a diuretic drug should be compared with the findings reported in this study for X-irradiation exposure.

#### VII. REFERENCES

1. Ulmer, D. D., L. B. Perkins, and J. G. Kereiakes. Alterations in Iodine-131 distribution in the rat after whole-body X-irradiation. USAMRL Report No. 345, Fort Knox, Ky, 1958; Rad. Res. 11: 810, 1959.
2. Donaldson, H. H. The Rat. Second edition, revised, Wistar Institute of Anatomy and Biology. Philadelphia, 1924.
3. Walker, W. G. and W. S. Wilde. Kinetics of radiopotassium in the circulation. Am. J. Physiol. 170: 401, 1952.
4. Hamilton, J. G. The metabolism of the fission products and the heaviest elements. Radiology, 49: 325, 1947.
5. Hood, S. L. and C. L. Comar. Metabolism of cesium-137 in laboratory and domestic animals. ORO-91, Univ. of Tenn., AEC Agricultural Research Program, 1953.

6. Hood, S. L. and C. L. Comar. Metabolism of cesium-137 in rats and farm animals. Arch. Biochem. and Biophys. 45: 423, 1953.
7. Ballou, J. E. and R. C. Thompson. Metabolism of cesium-137 in the rat: Comparison of acute and chronic administration experiments. Health Physics, 1: 85, 1958.
8. Edelman, A. Post-radiation water metabolism. AAAS . Symposium on Radiobiology. Nucleonics, 8(4): 28, 1951.
9. Edelman, A. and W. J. Eversole. Changes in antidiuretic activity of rat serum after X-irradiation. Am. J. Physiol. 163: 709, 1950.
10. Smith, W. W., I. B. Ackerman, and F. Smith. Body weight, fasting, and forced feeding after whole-body X-radiation. Am. J. Physiol. 168: 382, 1952.
11. Conrad, R. A. Some effects of ionizing radiation on the physiology of the gastrointestinal tract: A Review. Rad. Res. 5(2): 167, 1956.
12. Supplee, H., E. O. Weinman, and C. Entenman. Enlargement of the liver in Sprague-Dawley rats following whole-body X-irradiation. Am. J. Physiol. 185: 583, 1956.




Fig. 1. Small animal whole-body counting arrangement.




Fig. 2. Cesium-137 retention in starved control and X-irradiated rats.




Fig. 3. Urine excretion of cesium-137 (mean per cent of injected dose) in starved control and X-irradiated rats.




Fig. 4. Twenty-four hour distribution of cesium-137 in starved control and X-irradiated rats.

TABLE 1  
EFFECT OF STARVATION IN X-IRRADIATED RATS

Condition	Time Following Exposure	Body Weight	Feces Weight	Urine Volume	Water Intake
	days	grams	grams	ml	ml
Cesium-137 only	1	88.8 ± 1.9 (18)*	1.07 ± .54 (17)*	17 ± 7 (17)*	13 ± 8 (17)*
	2	82.9 ± 2.5 (18)	.42 ± .30 (17)	6 ± 3 (15)	7 ± 4 (15)
	3	77.7 ± 3.0 (18)	.32 ± .18 (17)	4 ± 3 (17)	4 ± 2 (9)
Cesium-137 and 300 r	1	88.9 ± 1.5 (9)	.31 ± .28 (9)	29 ± 11 (10)	22 ± 11 (10)
	2	82.7 ± 1.5 (10)	.31 ± .18 (9)	8 ± 6 (10)	9 ± 7 (10)
	3	76.8 ± 2.2 (10)	.76 ± .41 (10)	4 ± 4 (10)	3 ± 3 (10)
Cesium-137 and 600 r	1	91.0 ± 1.6 (13)	.39 ± .20 (12)	41 ± 15 (14)	45 ± 13 (13)
	2	83.2 ± 2.1 (14)	.34 ± .30 (12)	22 ± 12 (14)	18 ± 10 (12)
	3	77.2 ± 2.4 (14)	.54 ± .23 (14)	5 ± 4 (14)	6 ± 4 (14)
Cesium-137 and 900 r	1	88.2 ± 2.3 (19)	.46 ± .40 (17)	51 ± 19 (11)	44 ± 26 (16)
	2	80.4 ± 3.4 (19)	.38 ± .38 (14)	26 ± 11 (7)	26 ± 12 (18)
	3	74.3 ± 3.3 (18)	.40 ± .20 (17)	8 ± 3 (10)	7 ± 5 (19)

\*Numbers in parenthesis indicate number of rats.

TABLE 2  
CESIUM-137 RETENTION IN X-IRRADIATED RATS

Condition	Time Following Exposure	Total Body	Urine	Feces
	days	% of injected	% of injected	% of injected
Cesium-137 only	.007	94.5 ± 2.9 (18)*	.	.
	1	88.7 ± 1.9 (17)	6.8 ± 1.8 (16)*	0.8 ± 0.4 (12)*
	2	84.5 ± 2.8 (17)	2.6 ± 0.4 (13)	0.5 ± 0.3 (13)
	3	79.8 ± 2.9 (17)	2.5 ± 1.4 (16)	0.4 ± 0.2 (16)
Cesium-137 and 300 r	.007	94.8 ± 0.9 (8)	.	.
	1	87.3 ± 1.5 (9)	9.4 ± 1.3 (10)	0.2 ± 0.1 (8)
	2	83.7 ± 1.4 (9)	2.8 ± 0.8 (10)	0.4 ± 0.2 (10)
	3	78.2 ± 1.3 (8)	2.0 ± 0.8 (10)	0.6 ± 0.3 (9)
Cesium-137 and 600 r	.007	91.7 ± 1.8 (13)	.	.
	1	83.4 ± 1.9 (13)	11.7 ± 2.5 (14)	0.4 ± 0.3 (13)
	2	77.4 ± 2.9 (13)	4.3 ± 0.6 (13)	0.4 ± 0.2 (13)
	3	73.4 ± 2.9 (13)	2.6 ± 0.6 (13)	0.4 ± 0.2 (14)
Cesium-137 and 900 r	.007	93.8 ± 1.8 (18)	.	.
	1	87.1 ± 3.0 (18)	9.7 ± 2.3 (18)	0.4 ± 0.4 (18)
	2	80.0 ± 3.5 (18)	5.5 ± 1.1 (14)	0.3 ± 0.3 (14)
	3	74.5 ± 3.9 (18)	3.4 ± 0.8 (17)	0.4 ± 0.3 (18)

\*Numbers in parenthesis indicate number of rats.

TABLE 3  
CESIUM-137 DISTRIBUTION IN RAT TISSUES AT 24-HOURS POST-ADMINISTRATION

CUSTOM-137 DISTRIBUTION IN RAT TISSUES AT 24-HOURS POST-ADMINISTRATION

Organ or Body Fluid	Per Cent Dose per	Mean Per Cent of Dose Injected				Mean Concentration of Radioactivity							
		No. Rats	Control	No. Rats	X-irradiated	t	P	No. Rats	Control	No. Rats	X-irradiated	t	P
Blood	Total Volume	22	1.41 ± .23	21	1.23 ± .13	3.21	< 0.01	21	.26 ± .05	21	.23 ± .03	2.59	0.01
Pane	Organ	21	4.49 ± .72	20	2.94 ± .67	7.15	< 0.01	22	.71 ± .13	20	.44 ± .10	7.52	< 0.01
Gonads	Organ (both)	22	1.32 ± .28	20	1.26 ± .21	0.90	> 0.05	21	.95 ± .15	22	.93 ± .21	0.30	> 0.05
Heart	Organ	21	.67 ± .11	22	.67 ± .12	0.03	> 0.05	21	2.04 ± .30	21	1.96 ± .38	0.07	> 0.05
Kidney	Organ (both)	21	2.46 ± .55	21	2.37 ± .42	0.57	> 0.05	21	2.74 ± .55	22	2.66 ± .42	0.57	> 0.05
Large intestine	Organ & contents	22	3.96 ± 1.25	20	4.61 ± 2.17	1.18	> 0.05	20	1.44 ± .45	17	1.73 ± .71	1.44	> 0.05
Liver	Organ	22	6.24 ± 1.64	21	8.72 ± 1.38	5.48	< 0.01	22	1.76 ± .28	22	2.04 ± .43	2.58	0.01
Muscle	Organ	22	59.60 ± 9.03	22	53.80 ± 8.46	2.21	0.04	22	1.32 ± .21	22	1.20 ± .19	2.15	0.04
Small Intestine	Organ & contents	21	7.69 ± 1.31	22	6.42 ± 1.39	3.10	< 0.01	22	2.35 ± .47	22	2.11 ± .45	1.74	> 0.05
Spleen	Organ	21	.44 ± .10	20	.25 ± .06	8.00	< 0.01	22	1.67 ± .41	21	1.91 ± .30	2.27	0.03
Stomach	Organ & contents	20	1.66 ± .38	22	2.34 ± 1.43	2.15	0.04	21	2.26 ± .74	18	1.20 ± .34	5.91	< 0.01
Urine	Total Volume	22	9.15 ± 2.83	22	13.77 ± 5.10	3.71	< 0.01						

**DISTRIBUTION LIST OF USAMRL REPORTS**  
**Project No. 6X64-14-001 Biological and Medical**  
**Aspects of Ionizing Radiation**

**No. of  
Copies**

**AGENCY - DEFENSE**

**10 Armed Services Technical Information Agency, Arlington Hall Sta. Arlington Hall, Virginia**

**AGENCY - US ARMY**

- 6 Army Attache, Box 79, Navy 100 Fleet Post Office, New York, N. Y. . Attn: John C. Cressler, Colonel, Asst. Army Attache (Medical)**
- 3 Armed Forces Institute of Pathology, 6825 16th Street N. W., Washington 25, D. C.**
- 1 Brooke Army Medical Center, Medical Field Service School - Library - Fort Sam Houston, Texas**
- 1 Brooke Army Medical Center, U. S. Army Surgical Research Unit Library, Fort Sam Houston, Texas**
- 1 Brooke General Hospital, Medical Library, Box 151, Fort Sam Houston, Texas.**
- 1 Brooke General Hospital, Radioisotope Clinic, Brooke Army Medical Center, Fort Sam Houston, Texas**
- 1 Chief of Chemical Corps, Department of the Army, Washington 26, D. C., Attn: OCCCmlO/SA**
- 1 Chief of Engineers, Department of the Army, Washington 25, D. C., Attn: ENQNB**
- 1 Chief of Ordnance, CRDTB, Research and Special Projects Section, Washington 25, D. C**
- 1 Chief of Research and Development, Department of the Army, Chief of Life Science Division, Room 3D-442, The Pentagon, Washington 25, D. C.**
- 1 Chief of Research and Development, Department of the Army, Army Research Office, Washington 25, D. C., Attn: Scientific Information Branch**
- 1 Fitzsimons General Hospital, Medical Technical Library, Denver 30, Colorado**
- 1 Fitzsimons General Hospital, US Army Medical Research and Nutrition Laboratory, Denver 30, Colorado**
- 1 Letterman General Hospital, Medical Library, San Francisco, California**
- 1 Picatinny Arsenal, Office of the Commanding Officer, Dover, New Jersey, Attn: ORDBB-VS3**
- 1 Quartermaster Food and Container Institute, Library Branch, 1819 West Pershing Road, Chicago 9, Illinois**
- 1 Quartermaster Research and Engineering Field Evaluation Agency, Fort Lee, Virginia, Attn: Technical Library**
- 1 Quartermaster Research and Engineering Center, Natick, Massachusetts, Attn: Technical Library**
- 1 Redstone Arsenal, Redstone Arsenal, Alabama, Attn. ORDDW-HI**
- 1 William Beaumont General Hospital, El Paso, Texas**
- 1 US Army Alaska, Office of the Surgeon, APO 949, Seattle, Washington**
- 2 US Army Chemical Research and Development Center, Director, Army Chemical Center, Maryland**
- 1 US Army Chemical Corps Biological Laboratories, Fort Detrick, Maryland, Attn: Documents Technical Library Branch**
- 1 US Army Chemical Research and Development Laboratories, Technical Library, Building 330, Army Chemical Center, Maryland, Attn: Librarian**
- 1 US Army Command and General Staff College Library Services Br. Ft. Leavenworth Kansas, Attn: Archives**
- 1 US Army Dispensary, Springfield Armory, Springfield, Massachusetts, Attn: Medical Administrative Officer**
- 1 US Army Engineer Research and Development Laboratories, Fort Belvoir, Virginia, Attn: Technical Documents Center**
- 1 US Army Environmental Hygiene Agency, Army Chemical Center, Maryland**
- 1 US Army Europe, Medical Division, Plans and Operations Branch, APO 403, New York, N. Y.**
- 1 US Army Europe, Medical Laboratory, Department of Microbiology, APO 180, U. S. Forces, New York, New York**
- 1 US Army Hospital, Medical Library, Office of the Commanding Officer, Fort Lee, Virginia**
- 1 US Army Hawaii, Commanding General, APO 967, San Francisco, California, Attn: Surgeon**

# US ARMY - CONTINUED

## No. of Copies

- 1 US Army, Commanding General, I Corps Group, APO 358, San Francisco, California, Attn: Surgeon
- 1 US Army, Commanding General, 7th Logistical Command, APO 612, San Francisco, California, Attn: Surgeon
- 1 US Army, Commanding General 8th United States Army, APO, 301, San Francisco, California Attn: Surgeon
- 1 US Army, Japan, Commanding General, APO 343, San Francisco, California, Attn: Surgeon
- 1 US Army Leadership, Human Research Unit, Post Office Box 787, Presidio of Monterey, California, Attn: Library
- 2 US Army Liaison Group, Project Michigan, Willow Run Laboratories, Ypsilanti, Michigan
- 1 US Army Medical Command Japan, Medical General Laboratory (406) APO 343, San Francisco, California, Attn: Colonel Carl F. Tessmer, MC, Commanding
- 2 US Army Medical Liaison Branch, Office of the Chief Surgeon, Gorgas Hospital, Balboa Heights, Canal Zone
- 3 US Army Medical Research and Development Command, Commanding General, Main Navy Building, Washington 25, D. C.
- 1 US Army Medical Research Unit, Europe, Commanding Officer, APO 180, New York, New York
- 1 US Army Medical Research Unit, Commanding Officer, Institute for Medical Research, Kuala Lumpur, Malaya
- 1 US Army Medical Research Unit, Commanding Officer, Panama Field, Fort Clayton, Canal Zone
- 1 US Army Ryukyu Islands, IX Corps, Commanding General, APO 331, San Francisco, California, Attn: Surgeon
- 1 US Army Ordnance Human Engineering Laboratories, Aberdeen Proving Ground, Maryland
- 1 US Army Ordnance Arsenal, Frankford, Philadelphia 37, Pennsylvania, Attn: ORDA-1734/65-1
- 2 US Army Ordnance Tank-Automotive Command, Detroit Arsenal, Center Line, Michigan
- 1 US Army Research Office, Office of Chief Research and Development Command, Washington 25, D. C., Attn: Scientific Information Branch
- 1 US Army Signal Research and Development Laboratories, Commanding Officer, Fort Monmouth, New Jersey, Attn: Sig FM/EL-AITE
- 1 US Army Standardization Group, Canada, Office of the Senior Standardization Representative, c/o US Army Attache, US Embassy, Ottawa, Ontario, Canada
- 1 US Army Transportation Research Command, Fort Eustis, Virginia, Attn: Research Reference Center
- 1 US Army Tropical Research Medical Laboratory, APO 851, New York, New York
- 1 US Continental Army Command, Medical Section, Fort Monroe, Virginia
- 1 Walter Reed Army Institute of Research Department of Atomic Casualties Studies, Walter Reed Army Medical Center, Washington 12, D. C.
- 3 Walter Reed Army Institute of Research, Director, Walter Reed Army Medical Center, Washington 12, D. C.
- 1 Walter Reed Army Institute of Research, Division of Neuropsychiatry, Office of the Director, Washington 12, D. C.
- 1 Walter Reed Army Institute of Research, US Army Prosthetics Research Laboratory, Commanding Officer, Walter Reed Army Medical Center, Washington 12, D. C.
- 1 Walter Reed Army Medical Center, US Army Medical Service, Historical Unit, Washington 12, D. C.

## AGENCY - US NAVY

- 2 Bureau of Medicine and Surgery, Director, Research Division, Department of the Navy, Washington 25, D. C.
- 3 Bureau of Naval Weapons (DLI- 31) Department of the Navy, Washington 25, D. C.

No. of  
Copies

US NAVY - CONTINUED

- 2 Bureau of Yards and Docks, Department of the Navy, Washington 25, D. C., Attn: Code D-440
- 1 Chief of Naval Operations (OP03EG) Operations Evaluation Group, Department of the Navy, Washington 25, D. C.
- 1 Naval Medical Research Institute, Technical Reference Library, National Naval Medical Center, Bethesda, 14, Maryland
- 1 Office of Naval Research (Code 408), Department of the Navy, Special Assistant for Medical and Allied Sciences, Washington 25, D. C.
- 10 Office of Naval Research Branch Office, Navy 100, Box 39, Fleet Post Office, New York, New York, Attn: Victoria S. Greswold, Head, Administrative and Technical Service Br.
- 1 US Naval Air Development Center, Simulation Branch, Aeronautical Instrument Laboratory, Johnsville, Pennsylvania
- 1 US Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pennsylvania, Attn: Library
- 1 US Naval Civil Engineering Laboratory, Commanding Officer and Director (Code L31) Port Hueneme, California
- 2 US Naval Missile Center, Commander, Point Mugu, California, Attn: Technical Library
- 1 US Naval Medical School, National Naval Medical Center, Bethesda, 14, Maryland
- 1 US Naval Medical Research Laboratory, Box 100, US Naval Submarine Base, New London, Conn., Attn: Technical Library, Code 5111
- 1 US Naval Ordnance Test Station, The Medical Officer (Code 88), Station Hospital, Chica Lake, California
- 1 US Naval Radiological Defense Laboratory, Commanding Officer and Director (222), San Francisco 24, California
- 2 US Naval School of Aviation Medicine, US Naval Aviation Medical Center - 54, Pensacola, Florida
- 1 US Naval Supply Research and Development Facility, Clothing and Textile Division, 3rd Avenue and 29th Street, Brooklyn 32, New York, Attn: Library
- 2 US Naval Training Device Center, Human Engineering Department, Code 34, Port Washington, New York
- 1 US Naval Weapons Plant, Experimental Diving Unit, Washington 25, D. C.

AGENCY - AIR FORCE

- 1 Air Force Flight Test Center, Human Factors Branch (FTFH) Edwards Air Force Base, California
- AIR RESEARCH AND DEVELOPMENT COMMAND
- 1 Air Research and Development Command, US Air Force (RDGBL) Andrews Air Force Base, Washington 25, D. C.
  - 2 Air Research and Development Command, Rome Air Development Center, US Air Force, Griffiss Air Force Base, New York
  - 1 Air Development Division, US Air Force, Wright-Patterson Air Force Base, Ohio, Attn: WWRDAS Library
  - 1 Air Training Command (ATCSG-P) Randolph Air Force Base, Texas
  - 1 Arctic Aeromedical Laboratory, Commander, APO 731, Seattle, Washington, Attn: Librarian
  - 1 Assistant for Ground Safety, DCS/P, Headquarters US Air Force, Washington 25 D. C.
  - 1 Director of Research and Technology, Headquarters US Air Force, Washington 25, D. C., Attn: AFDRT-HF
  - 7 Ent Air Force Base, US Army Air Defense Command, Colorado Springs, Colorado, Attn: Colonel S. J. Neeson, MC, Command Surgeon
  - 3 Langley Research Center, National Aeronautics and Space Administration, Langley Field, Virginia, Attn: Librarian
  - 1 US Air Force, Strategic Air Command, (SUP3) Offutt Air Force Base, Nebraska
  - 2 US Air Force, SAM (SAMSGH-P) Brooks Air Force Base, Texas
  - 1 US Air Force Aerospace Medical Center (ATC), USAF Hospital, Lackland Air Force Base, Texas
  - 1 Wright Air Development Division, Wright-Patterson Air Force Base, Ohio, Attn: WWRDAS Aerospace Medical Library

No. of  
Copies

GOVERNMENTAL - AGENCIES

- 1 Argonne National Laboratory, 9700 South Cass Avenue, Argonne, Illinois, Attn: Hoylande D. Young
- 1 Central Intelligence Agency, 2430 E. Street, N. W., Washington, D. C., Attn: 1331, R&S Building
- 1 Chief, Defense Atomic Support Agency, Washington 25, D. C., Attn: Document Library Branch
- 1 Civil Aerosomedical Research Institute, Federal Aviation Agency, Post Office Box 1082, Oklahoma City, Oklahoma
- 1 Civil Defense Mobilization, Technical Reference Library, Battle Creek, Michigan, Attn: Virginia A. Staggers, Director
- 2 National Institute of Health, National Cancer Institute, Radiation Branch, Bethesda 14, Maryland
- 1 National Institute of Health, Library, Building 10 Room 5N118, Bethesda 14, Maryland, Attn: Acquisitions Section
- 1 National Institutes of Health, Division of Research Grants, Information Office, Bethesda, 14, Maryland
- 1 National Library of Medicine, Washington 25, D. C., Attn: Head Acquisition Section
- 1 National Research Council, Division of Medical Sciences, Medical Records 2101, Constitution Avenue, N. W., Washington 25, D. C.
- 2 US Atomic Energy Commission, Division of Biology and Medicine, Washington 23, D. C., Attn: Robert L. Corshie
- 1 US Atomic Energy Commission, Office of Technical Information Extension, Post Office Box 62, Oak Ridge, Tennessee

OTHER AGENCIES

- 1 Arctic Health Research Center, Library, Box 960, Anchorage, Alaska
- 2 Boeing Airplane Company, Aerospace Division, Seattle 23, Washington, Attn: Dr. Roney H. Lowry, Box 19-29, Chief, Space Medicine Section
- 1 Boeing Airplane Company, Library, Wichita Division, Wichita 1, Kansas
- 1 Boeing Airplane Company, Central Medical Library, Box 11-40, Post Office Box 3707, Seattle 24, Washington
- 1 Brookhaven National Laboratory, Research Library, Upton, Long Island, New York
- 2 Division of Radiological Health - ESS - Department of Health Education and Welfare, Room 5629 South HW Building, Washington 25, D. C.
- 1 General Electric Company, Advanced Electronics Center at Cornell University, Ithaca, New York, Attn: Library
- 1 General Electric Company, Technical Military Planning Operation, TEMPO Library, 735 State Street, Santa Barbara, California
- 1 John Crerar Library, 86 East Randolph Street, Chicago 1, Illinois
- 2 Kings County Hospital, Department of Anesthesiology, Brooklyn, New York, Attn: Dr. S. W. Weitzner
- 1 Lankenau Hospital, Division of Research, Lancaster and City Line Avenues, Philadelphia 31, Pennsylvania
- 1 Linfield Research Institute, McMinnville, Oregon
- 1 Mayo Clinic, Section of Biophysics, Rochester, Minnesota, Attn: Dr. Kenneth N. Ogle
- 1 Mercy Hospital, Anesthesia Research Laboratory, Pittsburgh 19, Pennsylvania
- 1 Motorola Incorporated, Systems Research Laboratory, 9330 Indiana Avenue, Riverside, California
- 7 National Aeronautics and Space Administration, 1510 H. Street, N. W., Washington 25, S. C., Attn: Bertram A. Mulcahy, Asst. Director for Technical Information
- 1 Rand Corporation, 1700 Main Street, Santa Monica, California, Attn: Library
- 1 Systems Research Center, Lockheed Electronics Company, Post Office Box 37, Bedminster, New Jersey
- 1 Yerkes Laboratories of Primate Biology, Incorporated, Orange Park, Florida, Attn: Dr. A. J. Riojelle

# MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

No. of  
Copies

- 1 Albany Medical College Library, New Scotland Avenue, Albany 8, New York
- 1 Bowman Gray School of Medicine Library, Winston-Salem, North Carolina
- 1 College of Medical Evangelists, Vernier Radcliffe Memorial Laboratory, Loma Linda, California
- 1 College of Medical Evangelists, White Memorial Medical Library, 1720 Brooklyn Avenue, Los Angeles 33, California
- 1 College of Physicians of Philadelphia, Library, 19 South 22nd Street, Philadelphia 3, Pennsylvania
- 1 Columbia University Medical Library, 639 West 168th Street, New York 32, New York
- 1 Cornell University Medical College Library, 1300 York Avenue, New York 21, New York
- 1 Creighton University, Medical Pharmacy Library, 1401 Devenport, Omaha 2, Nebraska
- 1 Dartmouth College Medical Library, Baker Building, Hanover, New Hampshire
- 1 Florida State University, Department of Psychology, Tallahassee, Florida, Attn: Dr. F. F. Dawson
- 1 Harvard Medical Library, 25 Shattuck Street, Boston 15, Massachusetts, Attn: Library
- 1 Indiana University Medical Center, School of Medicine Library, 1100 West Michigan Street, Indianapolis 7, Indiana
- 1 Indiana University Medical Center, 1100 West Michigan Street, Indianapolis 7, Indiana, Attn: Dr. Harris B. Shumacker, Jr., Professor of Surgery
- 1 Jefferson Medical College Library, 1025 Walnut Street, Philadelphia 7, Pennsylvania
- 1 Johns Hopkins University, Welch Medical Library, 1900 E. Monument Street, Baltimore 5, Maryland
- 1 Johns Hopkins University, Operations Research Office, Bethesda 14, Maryland, Attn: Library
- 1 Marquette University, Medical-Dental Library, 500 North 16th Street, Milwaukee 3, Wisconsin
- 1 Medical College of Virginia, Tompkins-McCaw Library, Richmond 19, Virginia
- 1 New York Academy of Medicine, Library, 2 East 103rd Street, New York, 29, New York
- 1 New York University Medical Center, Medical Library, 550 First Avenue, New York 18, New York
- 1 Northwestern University, Medical School, Archibald Church Library, 303 E. Chicago Avenue, Chicago 11, Illinois
- 1 Ohio State University, The Chemical Abstracts Service, Columbus 10, Ohio
- 1 Ohio State University, Engineering Experiment Station, The Systems Research Group, 156 W. 19th Avenue, Columbus 10, Ohio
- 1 Ohio State University, School of Optometry, Columbus 10, Ohio
- 1 Rush Medical College Library, 1758 West Harrison Street, Chicago 12, Illinois
- 1 Stanford University, Lane Medical Library, 300 Pasteur Road, Palo Alto, California
- 1 St. Louis University, Medical School Library, 1402 South Grand Boulevard, St. Louis 8, Missouri
- 1 State University of Iowa, College of Medicine Library, Medical Laboratories Building, Iowa City, Iowa
- 1 State University of New York, Downstate Medical Center, 450 Clarkson Avenue, Brooklyn 3, New York, Attn: Medical Librarian
- 1 Texas University Medical Center Library, Jesse H. Jones Library Building, Houston 25, Texas
- 1 Tulane University School of Medicine, 1430 Tulane Avenue, New Orleans 12, Louisiana, Attn: G. E. Burch, M.D., Professor of Medicine

# MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

No. of  
Copies

- 1 Vanderbilt University, School of Medicine, Nashville 5, Tennessee. Attn: George R. Keneely, M. D., Director, Radioisotope Center
- 2 West Virginia University, Medical Center Library, Morgantown, West Virginia
- 1 University of Alabama, Medical Center Library, 1919 Seventh Avenue South, Birmingham 3, Alabama
- 1 University of Arkansas, Medical Center Library, 4301 West Markham, Little Rock, Arkansas
- 1 University of Buffalo, Health Sciences Library, Copen Hall 141, Buffalo 14, New York
- 1 University of Buffalo, Psychology Department, Buffalo, New York
- 1 University of California, Medical Center, Biomedical Library, Los Angeles 24, California
- 1 University of California, 1301 South 46th Street, Richmond 4, California. Attn: Civil Defense Research Project
- 1 University of Chicago, US Air Force Radiation Laboratory, 930 E. 58th Street, Chicago 37, Illinois Attn: Director
- 1 University of Cincinnati, Kettering Laboratory, Eden and Bethesda Avenues, Cincinnati 19, Ohio
- 1 University of Florida, College of Medicine, Department of Physiology, Gainesville, Florida Attn: Dr. Melvin J. Fregly
- 2 University of Illinois, Aeromedical Laboratory, 840 South Wood Street, Chicago 12, Illinois
- 1 University of Illinois, Documents Division, Urbana, Illinois Attn: Librarian
- 1 University of Kansas Medical Center, Clendening Medical Library, Kansas City 12, Kansas
- 3 University of Louisville, School of Medicine Library, 101 W. Chestnut Street, Louisville 6, Kentucky
- 1 University of Maryland, Health Sciences Library, Serials Department, 111 South Greene Street, Baltimore 1, Maryland
- 1 University of Michigan, Serials and Documents Section, General Library, Ann Arbor, Michigan
- 1 University of Minnesota Library, Serials Division, Minneapolis 14, Minnesota
- 1 University of Missouri, Medical Library, Room M210 Medical Sciences Building, Columbia, Missouri
- 1 University of Nebraska, College of Medicine Library, 42nd and Dewey Avenue, Omaha 3, Nebraska
- 1 University of North Carolina, Memorial Hospital, Division of Health Affairs Library, Chapel Hill, North Carolina
- 1 University of Oklahoma Medical Center, Library, 801 N. E. 13th Street, Oklahoma City 4, Oklahoma
- 1 University of Oregon, Medical School Library, Portland 1, Oregon. Attn: Librarian
- 1 University of Pittsburgh, Falk Library of the Health Professions, Pittsburgh 13, Pennsylvania
- 1 University of Rochester, Atomic Energy Project, Technical Report Control Unit, Post Office Box 287, Station 3, Rochester 20, New York
- 1 University of Rochester, Strong Memorial Hospital, 260 Crittenden Boulevard, Rochester 20, New York
- 3 University of Southern California, School of Medicine Library, 2025 Zonal Avenue, Los Angeles 33, California
- 2 University of Tennessee, College of Medicine, Clinical Physiology, Institute of Clinical Investigation, 62, South Dunlap, Memphis 3, Tennessee
- 1 University of Tennessee Medical Units, Mooney Memorial Library, 62 South Dunlap, Memphis, 3, Tennessee, Attn: Librarian
- 1 University of Tennessee, Department of Bacteriology, Knoxville, Tennessee. Attn: Dr. D. Frank Holtman
- 1 University of Texas Medical Branch Library, Galveston, Texas. Attn: Librarian
- 1 University of Utah, Medical Library, Salt Lake City 12, Utah. Attn: Librarian
- 1 University of Vermont, College of Medicine Library, Burlington, Vermont

No. of  
Copies

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

- 1 University of Virginia Medical Library, University Hospital, Charlottesville, Virginia, Attn: Librarian
- 1 University of Washington, Health Sciences Library, Seattle 5, Washington
- 1 University of Wisconsin Medical School Library, SMI Building, N. Charter St. Madison 6, Wisconsin

FOREIGN

- 1 American Embassy, Office of the United States Army Attache, Foreign Service of the United States of America, London, England, Attn: Assistant Army Attache (Medical)
- 5 British Army Medical Liaison Officer, British Army Staff, British Embassy, Washington 8, D. C.
- 1 British Navy Staff Office, Benjamin Franklin Station, Post Office Box 165, Washington, D. C., Attn: Surgeon Captain F. P. Ellis
- 10 Canadian Liaison Office, US Army, Office of The Surgeon General, Room 1709A, Main Navy Building, Washington 25, D. C.
- 2 Defence Research Member, Canadian Joint Staff, 1450 Massachusetts Avenue, N. W., Washington 8, D. C.
- 1 German Military Attache, Federal Ministry of Defense, Washington 25, D. C.
- 1 Laboratory of General Pathology and Therapy, Professor Z. M. Bacq, 32, Bd. de la Constitution, Liege, Belgium
- 1 Royal Society of Medicine, Library, 1, Wimpole Street, London, W.I., England
- 1 University of Alberta, University Medical Library, Edmonton, Alberta, Canada, Attn: Phyllis Russell, Medical Librarian
- 2 University of Western Ontario, Medical School, Department of Biophysics, South Street, London, Ontario, Canada, Attn: Dr. Allen C. Burton

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
CESIUM-137 RETENTION AND DISTRIBUTION IN X-  
IRRADIATED RATS - J. G. Kerelakes, D. D.  
Uimer, A. T. Kreba, and T. D. Sterling  
Report No. 504, 4 Sep 61, 12 pp & ii, 4  
illus. - 3 tables - Project No. 6X04-14-001,  
Unclassified Report

UNCLASSIFIED  
1. Radiobiology  
2. Isotope  
Metabolism  
3. Cesium-137

The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1  $\mu$ C cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5  $\mu$ C cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
CESIUM-137 RETENTION AND DISTRIBUTION IN X-  
IRRADIATED RATS - J. G. Kerelakes, D. D.  
Uimer, A. T. Kreba, and T. D. Sterling  
Report No. 504, 4 Sep 61, 12 pp & ii, 4  
illus. - 3 tables - Project No. 6X04-14-001,  
Unclassified Report

UNCLASSIFIED  
1. Radiobiology  
2. Isotope  
Metabolism  
3. Cesium-137

The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1  $\mu$ C cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5  $\mu$ C cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
CESIUM-137 RETENTION AND DISTRIBUTION IN X-  
IRRADIATED RATS - J. G. Kerelakes, D. D.  
Uimer, A. T. Kreba, and T. D. Sterling  
Report No. 504, 4 Sep 61, 12 pp & ii, 4  
illus. - 3 tables - Project No. 6X04-14-001,  
Unclassified Report

UNCLASSIFIED  
1. Radiobiology  
2. Isotope  
Metabolism  
3. Cesium-137

The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1  $\mu$ C cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5  $\mu$ C cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
CESIUM-137 RETENTION AND DISTRIBUTION IN X-  
IRRADIATED RATS - J. G. Kerelakes, D. D.  
Uimer, A. T. Kreba, and T. D. Sterling  
Report No. 504, 4 Sep 61, 12 pp & ii, 4  
illus. - 3 tables - Project No. 6X04-14-001,  
Unclassified Report

UNCLASSIFIED  
1. Radiobiology  
2. Isotope  
Metabolism  
3. Cesium-137

The influence of external ionizing radiation on the retention and distribution of cesium-137 was studied. Immediately following a total-body exposure of 300, 600, or 900 r of 250 kv X-rays, rats were injected with 1  $\mu$ C cesium-137. Rats given cesium-137 only served as controls. The X-irradiated rats excreted more cesium-137 over the three-day period following the irradiation than the controls, primarily accounted for by increasing urine excretion. 900 r X-irradiated rats injected with 5  $\mu$ C cesium, sacrificed at 24 hours, retained less cesium in muscle, bone, blood, small intestine, and spleen but more in stomach and liver than the controls. Possible mechanisms are discussed.